

Seed Germination of Alaska Iris, *Iris setosa* ssp. *interior*

Patricia S. Holloway¹

School of Agriculture and Land Resources Management, University of Alaska, Fairbanks, AK 99775-0100

Additional index words. dormancy, stratification, gibberellic acid, light

Abstract. Seeds of Alaska iris, *Iris setosa* Pall. ssp. *interior* (Anders.) Hult., were collected from wild stands near Fairbanks and subjected to the following treatments: 0 or 125 days of stratification; 24-hr soak in water or 1000 mg·liter⁻¹ gibberellic acid (GA₃); alternating (25°/10°C) or constant (21°) temperatures; and germination in darkness or light. After 7 days, germination was best (95.0%) with stratified seeds that were soaked in GA₃ and germinated at constant temperatures in the dark. A greenhouse study in which stratified seeds were soaked in water or GA₃, sown in a commercial peat and vermiculite seed germination mix, and germinated beneath clear or black plastic confirmed that germination was highest (64.4%) and most uniform (83.3% filled container cells) with the combined treatments of GA₃ and darkness.

Seeds of Alaska iris do not germinate immediately upon harvest. Seeds require a period of cold stratification ranging from 120 to 150 days at 4°C, but germination rarely exceeds 40%. This germination pattern is similar to other *Iris* species and hybrids in which dormancy may persist for several months, followed by intermittent germination over several years. (2, 3, 9, 11, 12, 15). Dormancy is reported to be caused by a dormant embryo (9), a water-soluble inhibitor in the endosperm (3, 4, 8, 12, 14), and/or mechanical inhibition of embryo growth (5, 14).

Several methods have been used to improve germination percentages of iris seeds, including cold stratification (9, 10, 12), leaching (3, 12), chipping or removal of the seedcoat (5, 12, 15); sowing immature seeds (2, 6), heat treatment (7), and embryo excision (9, 11-13). With the exception of embryo excision, these methods have not consistently improved germination percentages. None of these treatments has been reported previously with the Alaska iris to my knowledge. The purpose of this experiment was to test various treatments including cold stratification, light, gibberellic acid, and temperature to determine possible methods of improving germination of the Alaska iris.

Seed capsules of Alaska iris were collected on 22 Aug. 1985 from wild stands growing near Fairbanks. Capsules were air-dried for 24 hr to facilitate removal of the seeds; seeds were then stored dry at 4°C for 2 weeks. One-half of the seeds were stratified in peat at 4° for 125 days, while the remainder were stored dry at 4° until further

treatment. Both stratified and unstratified seeds were soaked in tap water (pH 6.5) or 1000 mg·liter⁻¹ gibberellic acid (GA₃) for 24 hr at room temperature, rinsed in water, and then placed on moistened filter paper in petri dishes. One-half of the dishes were enclosed in aluminum foil to exclude light, and all dishes were placed into clear plastic bags to retard evaporation. Dishes were placed in a growth chamber at 21° constant temperature. Light was provided for 16 hr each day using a mixture of 40-W cool-white fluorescent and 25-W incandescent lamps with a radiant flux of 55 W·m⁻² measured 33 cm below the lamps using a LI-COR Model LI-185B Quantum/Radiometer/Photometer. Previous experiments had shown that non-dormant seeds germinate rapidly in 3 to 5 days with little or no additional germination after 7 days. Consequently, germination was recorded as radicle emergence from the seedcoat after 7 days of treatment. The entire experiment was repeated under alternating temperatures of 25° (day) and 10° (night) for 16 and 8 hr, respectively. Both experiments were conducted as a completely randomized design with three replicates per treatment and 50 seeds per replicate.

In order to test selected treatments under greenhouse conditions, seeds from a different wild stand near Fairbanks were collected on 1 Sept. 1985. Seeds were stratified in peat at 4°C for 125 days and then soaked in 1000 mg·liter⁻¹ GA₃ or water. Three seeds per cell were sown ≈5 mm deep in six-celled plastic containers filled with a commercial peat and vermiculite seed germination mix (≈45 cm³ of mix per cell). Containers were watered and enclosed in individual black or clear plastic bags, which then were placed randomly on a bench in the greenhouse. A minimum air temperature of 21° was maintained. Thermocouples were inserted at random beneath the plastic to a medium depth of 5 mm in order to determine any temperature differences between plastics. Since medium temperatures did not differ by more than 3° between plastic treatments, temper-

Table 1. Effects of light, GA₃, and constant or alternating temperatures on seed germination of Alaska iris in a growth chamber.

Treatment	Germination (%) ²			
	21°C		25°/10°	
	Mean	SE	Mean	SE
Light				
GA ₃ soak	53.6	2.4	23.3	6.2
Water soak	33.3	6.2	25.0	8.2
Dark				
GA ₃ soak	95.0	8.5	35.0	4.1
Water soak	30.0	7.1	35.0	4.1
Significance				
Light/dark	*		NS	
GA ₃ /water	**		NS	
Treatment interaction	**		NS	
GA ₃ /water × Light	*			
GA ₃ /water × Dark	**			
Light/Dark × GA ₃	**			
Light/Dark × Water	NS			

²Radicle emergence after 7 days.

NS,*,**Not significant and significant at P = 5% and 1%, respectively.

Table 2. Effects of GA₃ or water soak and clear or black plastic on germination of Alaska iris seeds sown in a peat and vermiculite mix.

Treatment	Seedling emergence (%) ²		Empty container cells (%)	
	Mean	SE	Mean	SE
Black plastic covering				
GA ₃ soak	64.4	13.8	16.7	11.5
Water soak	50.5	12.1	30.6	14.9
Clear plastic covering				
GA ₃ soak	57.0	13.6	29.7	13.7
Water soak	38.1	10.2	59.8	15.9
Average treatment effects				
Between plastics				
Black	57.5**		23.6**	
Clear	48.0		44.4	
Between water and GA ₃ soak				
GA ₃ soak	60.7**		22.9**	
Water soak	44.2		45.1	

²Shoot emergence from medium after 30 days.

**Treatment pairs within columns significant at 1%.

ature was not considered further as a major treatment effect. Sunlight was supplemented with a 16-hr photoperiod from 6:00 AM to 10:00 PM using 400-W high-pressure sodium lamps at 48 W·m⁻² measured to 1.6 m below the lamps. After 10 days, all containers were removed from the plastic bags. After an additional 30 days, the percent seedling emergence from the medium was recorded. Data were analyzed using analysis of variance for four factors with treatment containers replicated 12 times in a completely randomized design. Where appropriate, data were subject to arcsin transformation prior to analysis.

Unstratified seeds failed to germinate regardless of constant or alternating temperatures, GA₃ or light treatments, and consequently the data were omitted from further treatment comparisons. In a comparison of the stratified seeds, the GA₃ treatment significantly improved germination at constant temperatures in both light and dark (Ta-

Received for publication 17 Nov. 1986. Paper no. J-178 of the Scientific Journal Series of the Univ. of Alaska Agricultural and Forestry Experiment Station. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.

¹Assistant Professor.

ble 1). Germination also was improved by the dark treatment, but was the best with a combination of GA₃ and darkness. Differences between light and dark germination in water-soaked seeds were not significant.

Germination at alternating temperatures never exceeded 45% for any treatment replicate, and differences among treatments were not significant (Table 1). Although it appeared that a constant temperature was preferable to alternating temperatures, the 8 hr of low temperatures per day may have delayed germination so that maximum germination and treatment effects were not evident after the 7-day germination period.

Results of the greenhouse experiments were similar to the growth chamber tests in that GA₃ significantly improved seedling emergence of stratified seeds (Table 2). Emergence also was improved by covering the flats with black plastic, but there was no significant interaction between the two main treatments. Since all seeds were sown at an ≈5-mm depth, all seeds in flats covered with either clear or black plastic received a reduced light treatment, which could explain the lack of interaction evident in the previous experiment and the similar germination percentages recorded for GA₃-treated seeds in both plastic treatments.

The uniformity of seedling emergence is reflected in the percentage of empty cells per container (Table 2). Stratified seeds that were soaked in water and germinated in the clear plastic bags showed no seedling emergence in more than half of the container cells. Based on the percentage of empty cells for this treatment, an average of 7.4 seeds per cell would have to be sown in order to approach 100% filled cells. In contrast, the stratified, GA₃-treated seeds that were germinated beneath black plastic had only 16.7% empty cells, and would require an average of 3.6 seeds per cell. The remaining two treatments would require an average of 4.3 seeds per cell.

Both experiments demonstrated that germination is improved significantly by soaking the stratified seeds in 1000 mg·liter⁻¹ GA₃ and germinating them in darkness. Reports of GA₃ treatments using the *Iris* spp. have been limited. Abdalla and McKelvie (1) reported that *I. mellita* Janka seeds failed to germinate when they were soaked in 500 mg·liter⁻¹ GA₃ and then held at 23°C. Seeds soaked in GA₃ and stratified at 1.5° for 4 weeks before being transferred to the higher temperature also did not germinate. The GA₃ treatment in both *I. setosa* and *I. mellita* apparently does not replace the chilling requirement of unstratified seeds. The 4-week chilling period probably was too short to overcome dormancy in the *I. mellita* seeds.

Few reports document the response of iris seeds to darkness. Guppy (7) found that 46.6% of *I. pseudacorus* L. seeds germinated after 6 months in the greenhouse in containers of water that were covered to exclude light. No seeds germinated in the light at room temperature or in the greenhouse, or in the dark at room temperature. Randolph (11) recommended culturing excised em-

bryos first in darkness at 28° to 30°C for 3 to 5 days, followed by a gradual increase in light until seedlings were transferred to a shaded greenhouse. Both of these studies indicate a positive effect of darkness on germination. The experiments with *I. setosa* showed that darkness does not influence germination of unstratified seeds nor is it an obligate requirement of stratified seeds. Darkness does improve germination percentages, particularly when combined with GA₃ treatment. This combination of stratification, GA₃ and darkness provides a less cumbersome method than embryo excision for obtaining high germination percentages for commercial producers and plant breeders.

Literature Cited

1. Abdalla, S.T. and A.D. McKelvie. 1980. The interaction of chilling and gibberellic acid on the germination of seeds of ornamental plants. *Seed Sci. & Technol.* 8:139-144.
2. Arceneaux, G. 1941. Breeding and propagating Louisiana iris. *Bul. Amer. Iris. Soc.* 82:67-71.
3. Arditti, J. and T.R. Pray. 1969. Dormancy factors in iris (Iridaceae) seeds. *Amer. J. Bot.* 56:254-259.
4. Cox, L.G. 1942. A physiological study of embryo dormancy in the seed of native hard-

woods and iris. PhD Diss., Cornell Univ., Ithaca, N.Y.

5. Crocker, W. 1906. Role of seed coats in delayed germination. *Bot. Gaz.* 42:265-92.
6. Essig, E.O. 1934. Twelve years of intensive iris breeding in California, methods and records. *Bul. Amer. Iris. Soc.* 52:3-30.
7. Guppy, H.B. 1897. On the postponement of the germination of the seeds of aquatic plants. *Proc. R. Phys. Soc. Sess. 1896-97.* 13:344-359.
8. Jorgensen, C.J.C. 1965. Inhibitory effects of iris seed extracts on germination of indicator plants. *Bul. Amer. Iris Soc.* 179:27-33.
9. Lenz, L.W. 1955. Studies in iris embryo culture I. Germination of embryos of the subsection *Hexapogon* Benth (Sect. *Regelia* sensu Dykes). *El Aliso* 3:173-182.
10. Meserve, J.C. 1985. Propagation of iris 'Pacific Coast Hybrids'. *The Plant Prop.* 31:6.
11. Randolph, L.F. 1945. Embryo culture of iris seeds. *Bul. Amer. Iris Soc.* 97:33-45.
12. Randolph, L.F. and L.G. Cox. 1943. Factors influencing the germination of iris seed and the relation of inhibiting substances to embryo dormancy. *Proc. Amer. Soc. Hort. Sci.* 43:284-300.
13. Slatter, E.M. 1950. The culture of iris embryos on nutrient agar. *Bul. Kew R. Bot. Garden* 3:425-430.
14. Stoltz, L.P. 1968. Iris seed dormancy. *Physiol. Plant.* 21:1328-1331.
15. White, C.G. 1936. Experiments with seed germination. *Bul. Amer. Iris Soc.* 97:46.